

The Geriatric Depression Scale (GDS)

By: Sherry A. Greenberg, PhD(c), MSN, GNP-BC,
Hartford Institute for Geriatric Nursing, NYU College of Nursing

WHY: Depression is common in late life, affecting nearly 5 million of the 31 million Americans aged 65 and older with clinically significant depressive symptoms reaching 13% in older adults aged 80 and older (Blazer, 2009). Major depression is reported in 8-16% of community dwelling older adults, 5-10% of older medical outpatients seeing a primary care provider, 10-12% of medical-surgical hospitalized older adults with 23% more experiencing significant depressive symptoms (Blazer, 2009). Recognition in long-term care facilities is poor and not consistent amongst studies (Blazer, 2009).

Depression is not a natural part of aging. Depression is often reversible with prompt recognition and appropriate treatment. However, if left untreated, depression may result in the onset of physical, cognitive, functional, and social impairment, as well as decreased quality of life, delayed recovery from medical illness and surgery, increased health care utilization, and suicide.

BEST TOOL: While there are many instruments available to measure depression, the Geriatric Depression Scale (GDS), first created by Yesavage, et al., has been tested and used extensively with the older population. The GDS Long Form is a brief, 30-item questionnaire in which participants are asked to respond by answering yes or no in reference to how they felt over the past week. A Short Form GDS consisting of 15 questions was developed in 1986. Questions from the Long Form GDS which had the highest correlation with depressive symptoms in validation studies were selected for the short version. Of the 15 items, 10 indicated the presence of depression when answered positively, while the rest (question numbers 1, 5, 7, 11, 13) indicated depression when answered negatively. Scores of 0-4 are considered normal, depending on age, education, and complaints; 5-8 indicate mild depression; 9-11 indicate moderate depression; and 12-15 indicate severe depression.

The Short Form is more easily used by physically ill and mildly to moderately demented patients who have short attention spans and/or feel easily fatigued. It takes about 5 to 7 minutes to complete.

TARGET POPULATION: The GDS may be used with healthy, medically ill and mild to moderately cognitively impaired older adults. It has been extensively used in community, acute and long-term care settings.

VALIDITY AND RELIABILITY: The GDS was found to have a 92% sensitivity and a 89% specificity when evaluated against diagnostic criteria. The validity and reliability of the tool have been supported through both clinical practice and research. In a validation study comparing the Long and Short Forms of the GDS for self-rating of symptoms of depression, both were successful in differentiating depressed from non-depressed adults with a high correlation ($r = .84, p < .001$) (Sheikh & Yesavage, 1986).

STRENGTHS AND LIMITATIONS: The GDS is not a substitute for a diagnostic interview by mental health professionals. It is a useful screening tool in the clinical setting to facilitate assessment of depression in older adults especially when baseline measurements are compared to subsequent scores. It does not assess for suicidality.

FOLLOW-UP: The presence of depression warrants prompt intervention and treatment. The GDS may be used to monitor depression over time in all clinical settings. Any positive score above 5 on the GDS Short Form should prompt an in-depth psychological assessment and evaluation for suicidality.

MORE ON THE TOPIC:

Best practice information on care of older adults: www.ConsultGeriRN.org.

The Stanford/VA/NIA Aging Clinical Resource Center (ACRC) website. Retrieved July 2, 2012, from <http://www.stanford.edu/~yesavage/ACRC.html>. Information on the GDS. Retrieved July 2, 2012, from <http://www.stanford.edu/~yesavage/GDS.html>.

Blazer, D.G. (2009). Depression in late life: Review and commentary. *FOCUS*, 7(1), 118-136.

Greenberg, S.A. (2007). How to Try This: The Geriatric Depression Scale: Short Form. *AJN*, 107(10), 60-69.

Harvath, T.A., & McKenzie, G. (2012). Depression in Older Adults. In M. Boltz, E. Capezuti, T.T. Fulmer, & D. Zwicker (Eds.), A. O'Meara (Managing Ed.), *Evidence-based geriatric nursing protocols for best practice* (4th ed., pp. 135-162). NY: Springer Publishing Company, LLC.

Koenig, H.G., Meador, K.G., Cohen, J.J., & Blazer, D.G. (1988). Self-rated depression scales and screening for major depression in the older hospitalized patient with medical illness. *JAGS*, 36, 699-706.

Sheikh, J.I., & Yesavage, J.A. (1986). Geriatric Depression Scale (GDS). Recent evidence and development of a shorter version. In T.L. Brink (Ed.), *Clinical Gerontology: A Guide to Assessment and Intervention* (pp. 165-173). NY: The Haworth Press, Inc.

Yesavage, J.A., Brink, T.L., Rose, T.L., Lum, O., Huang, V., Adey, M.B., & Leirer, V.O. (1983). Development and validation of a geriatric depression screening scale: A preliminary report. *Journal of Psychiatric Research*, 17, 37-49.

Geriatric Depression Scale: Short Form

Choose the best answer for how you have felt over the past week:

1. Are you basically satisfied with your life? YES / **NO**
2. Have you dropped many of your activities and interests? **YES** / NO
3. Do you feel that your life is empty? **YES** / NO
4. Do you often get bored? **YES** / NO
5. Are you in good spirits most of the time? YES / **NO**
6. Are you afraid that something bad is going to happen to you? **YES** / NO
7. Do you feel happy most of the time? YES / **NO**
8. Do you often feel helpless? **YES** / NO
9. Do you prefer to stay at home, rather than going out and doing new things? **YES** / NO
10. Do you feel you have more problems with memory than most? **YES** / NO
11. Do you think it is wonderful to be alive now? YES / **NO**
12. Do you feel pretty worthless the way you are now? **YES** / NO
13. Do you feel full of energy? YES / **NO**
14. Do you feel that your situation is hopeless? **YES** / NO
15. Do you think that most people are better off than you are? **YES** / NO

Answers in **bold** indicate depression. Score 1 point for each bolded answer.

A score > 5 points is suggestive of depression.

A score ≥ 10 points is almost always indicative of depression.

A score > 5 points should warrant a follow-up comprehensive assessment.

Source: <http://www.stanford.edu/~yesavage/GDS.html>

This scale is in the public domain.

The Hartford Institute for Geriatric Nursing would like to acknowledge the original author of this Try This, Lenore Kurlowicz, PhD, RN, CS, FAAN, who made significant contributions to the field of geropsychiatric nursing and passed away in 2007.

Geriatric Depression Scale (short form)

Instructions: Circle the answer that best describes how you felt over the past week.

- | | | |
|---|-----|----|
| 1. Are you basically satisfied with your life? | yes | no |
| 2. Have you dropped many of your activities and interests? | yes | no |
| 3. Do you feel that your life is empty? | yes | no |
| 4. Do you often get bored? | yes | no |
| 5. Are you in good spirits most of the time? | yes | no |
| 6. Are you afraid that something bad is going to happen to you? | yes | no |
| 7. Do you feel happy most of the time? | yes | no |
| 8. Do you often feel helpless? | yes | no |
| 9. Do you prefer to stay at home, rather than going out and doing things? | yes | no |
| 10. Do you feel that you have more problems with memory than most? | yes | no |
| 11. Do you think it is wonderful to be alive now? | yes | no |
| 12. Do you feel worthless the way you are now? | yes | no |
| 13. Do you feel full of energy? | yes | no |
| 14. Do you feel that your situation is hopeless? | yes | no |
| 15. Do you think that most people are better off than you are? | yes | no |

Total Score _____

Geriatric Depression Scale (GDS)

Scoring Instructions

Instructions: Score 1 point for each bolded answer. A score of 5 or more suggests depression.

- | | | |
|---|------------|-----------|
| 1. Are you basically satisfied with your life? | yes | no |
| 2. Have you dropped many of your activities and interests? | yes | no |
| 3. Do you feel that your life is empty? | yes | no |
| 4. Do you often get bored? | yes | no |
| 5. Are you in good spirits most of the time? | yes | no |
| 6. Are you afraid that something bad is going to happen to you? | yes | no |
| 7. Do you feel happy most of the time? | yes | no |
| 8. Do you often feel helpless? | yes | no |
| 9. Do you prefer to stay at home, rather than going out and doing things? | yes | no |
| 10. Do you feel that you have more problems with memory than most? | yes | no |
| 11. Do you think it is wonderful to be alive now? | yes | no |
| 12. Do you feel worthless the way you are now? | yes | no |
| 13. Do you feel full of energy? | yes | no |
| 14. Do you feel that your situation is hopeless? | yes | no |
| 15. Do you think that most people are better off than you are? | yes | no |

A score of ≥ 5 suggests depression

Total Score _____

Ref. Yes average: The use of Rating Depression Series in the Elderly, in Poon (ed.): Clinical Memory Assessment of Older Adults, American Psychological Association, 1986